

Attachment 2

Concentrations of Germicides for General Environmental Sanitation Use

| Active Component Chinese/English Name | Application Type | Concentration |
|---|--------------------------------|---------------|
| Name of Active ingredient | | |
| Alkyl dimethyl benzyl ammonium saccharinate | Liquid, aerosol spray | 4.80 % |
| Benzalkonium chloride | Liquid, aerosol spray | 4.80 % |
| Didecylmethyl ammonium chloride | Liquid, aerosol spray | 1.68 % |
| Glutaraldehyde | Liquid, aerosol spray | 2.68 % |
| n-Alkyldimethylbenzyl ammonium chloride | Liquid, aerosol spray | 4.80 % |
| Obanol-516 | Liquid, aerosol spray | 5.00 % |
| o-Benzyl-p-chlorophenol | Liquid, aerosol spray | 5.00 % |
| o-Phenylphenol | Liquid, aerosol spray | 5.00 % |
| p-Chlorophenol | Liquid, aerosol spray | 5.00 % |
| p-Dichlorobenzene | Liquid, aerosol spray | 5.00 % |
| Polyalkyl polyamino ethylglycine | Liquid, aerosol spray | 5.00 % |
| p-ter-Amylphenol | Liquid, aerosol spray | 5.00 % |
| Calcium hypochlorite, $\text{Ca}(\text{ClO})_2$ | Pellet, tablet, powder, liquid | 90 % |
| Chlorine dioxide, ClO_2 | Liquid, powder, tablet | 50 % |
| Sodium chlorite, NaClO_2 | Liquid, powder, tablet | 50 % |
| Sodium hypochlorite, NaClO | Powder | 50 % |
| Bleach | Powder | 80 % |